



Diversity of wood – decaying fungi in Haikou Forestry Farm, Yunnan Province, P.R. China

He X, Chen JZ and Zhao CL*

College of Biodiversity Conservation, Southwest Forestry University, Kunming 650224, P.R. China

He X, Chen JZ, Zhao CL 2021 – Diversity of wood-decaying fungi in Haikou Forestry Farm, Yunnan Province, P.R. China. Studies in Fungi 6(1), 365–377, Doi 10.5943/sif/6/1/27

Abstract

Nine field trips carried out in Haikou Forestry Farm, Yunnan Province, P.R. China resulted in 681 specimens of wood-decaying fungi. The present paper summarizes 52 species collected that are distributed in 37 genera, 16 families, 6 orders including their hosts and substrates. A checklist of wood-decaying fungi in Haikou Forestry Farm is also given. Phylogenetic analysis of ITS nrRNA gene region was performed for all the collected samples with maximum likelihood, maximum parsimony and Bayesian inference methods. The phylogenetic tree showed that fifty-two species nested in sixteen families belonging to six orders in Agaricomycetes.

Keywords – Basidiomycota – biological resources – diversity – Haikou Forestry Farm – mycota

Introduction

The diversity for flora of seed plants in Yunnan Province, P.R. China is observably high, and the endemic species of woody plants are rich, in which both supply good substrates for wood-decaying fungi. Wood-decaying fungi are kind of large basidiomycetes that grow on various kinds of wood, such as the living trees, dead standing trees, fallen trunk, fallen branch and stump (Dai 2012a), which can be used for industrial value, medicinal value, edible value and economic value (Russell & Paterson 2006, Dai et al. 2015, Vinay et al. 2015, M'Barek et al. 2020, Wu et al. 2020, Runnel et al. 2021).

Haikou Forestry Farm is located in Haikou Town, Kunming, belonging to the Jinsha River system and the geographical location is between 102°28'-102°38' E and 24°4'-24°56' N, with the altitude of 1800-2400 m (Xu et al. 2015, Zhang et al. 2017). It has a subtropical monsoon climate and the native tree species with mostly subtropical evergreen broadleaf trees (Xiong & Zhou 2019). The main vegetation includes warm coniferous forest (*Pinus yunnanensis* Franch., *P. armandii* Franch. and *Keteleeria evelyniana* Mast.), deciduous broad-leaved forest (*Alnus nepalensis* D. Don), and semi-humid evergreen broad-leaved forest (*Lithocarpus dealbatus* (J. D. Hooker et Thomson ex Miquel) Rehder and *Castanopsis delavayi* Franch.).

However, the previously documented wood-decaying fungi are mostly in northwest Yunnan Province, China, and few polypore and corticioid fungi have been reported in here so far. According to the modern taxonomy (Dai 2012a), wood-decaying fungi mainly belong to ten orders of Agaricomycetes, viz., Agaricales, Auriculariales, Cantharellales, Corticiales, Gloeophyllales, Hymenochaetales, Polyporales, Russulales, Thelephorales and Trechisporales. Therefore, the current wood-decaying fungi catalogues include poroid and corticioid hymenophore. In the present study, nine field trips were carried out in different areas of Haikou Forestry Farm, and about 681 specimens were collected, in which 52 species belonging to 37 genera, 16 families, 6 orders, which

were identified from these materials. This paper is going to summarize the distribution of wood-decaying fungi and enrich the fungal diversity in this area.

Materials & Methods

Morphological studies

The studied specimens are deposited at the herbarium of Southwest Forestry University (SWFC), Kunming, Yunnan Province, P.R. China. Macromorphological descriptions were based on field notes. Colour terms are from Petersen (1996). Micromorphological data were obtained from the dried specimens, and observed under a light microscope following Dai (2012a). The following abbreviations were used for the micro characteristics description: KOH = 5% potassium hydroxide, CB = Cotton Blue, CB⁻ = acyanophilous, CB⁺ = cyanophilous, IKI = Melzer's reagent, IKI⁻ = both inamyloid and indextrinoid, L = mean spore length (arithmetic average of all spores), W = mean spore width (arithmetic average of all spores), Q = variation in the L/W ratios between the specimens studied, n (a/b) = number of spores (a) measured from given number (b) of specimens.

Molecular procedures and phylogenetic analysis

CTAB rapid plant genome extraction kit-DN14 (Aidlab Biotechnologies Co., Ltd, Beijing) was used to obtain genomic DNA from dried specimens, according to the manufacturer's instructions that a small piece of dried fungal specimen (about 30 mg) was ground to powder with liquid nitrogen. The powder was transferred to a 1.5 mL centrifuge tube, suspended in 0.4 mL of lysis buffer, and incubated in a 65°C water bath for 60 min. After that, 0.4 mL phenol-chloroform (24:1) was added to each tube and the suspension was shaken vigorously. After centrifugation at 13,000 rpm for 5 min, 0.3 mL supernatant was transferred to a new tube and mixed with 0.45 mL binding buffer. The mixture was then transferred to an adsorbing column (AC) for centrifugation at 13,000 rpm for 0.5 min. Then, 0.5 mL inhibitor removal fluid was added in AC for a centrifugation at 12,000 rpm for 0.5 min. After washing twice with 0.5 mL washing buffer, the AC was transferred to a clean centrifuge tube, and 100 mL elution buffer was added to the middle of adsorbed film to elute the genome DNA. ITS region was amplified with primer pair ITS5 and ITS4 (White et al. 1990). The PCR procedure for ITS was as follows: initial denaturation at 95°C for 3 min, followed by 35 cycles at 94°C for 40 s, 58°C for 45 s and 72°C for 1 min, and a final extension of 72°C for 10 min. The PCR products were purified and directly sequenced at Kunming Tsingke Biological Technology Limited Company, Kunming Yunnan Province, P.R. China. All newly generated sequences were deposited at GenBank (Table 1).

Sequencher 4.6 (GeneCodes, Ann Arbor, MI, USA) was used to edit the DNA sequence. Sequences were aligned in MAFFT 7 (<http://mafft.cbrc.jp/alignment/server/>) using the "G-INS-i" strategy and manually adjusted in BioEdit (Hall 1999). Sequences of *Tremella flava* Chee J. Chen and *T. mesenterica* Retz. obtained from GenBank was used as an outgroup to root tree following He et al. (2019) in ITS analysis (Fig. 1).

Maximum parsimony analysis was applied to the ITS dataset sequences. Approaches to phylogenetic analysis followed Zhao & Wu (2017) and the tree construction procedure was performed in PAUP* version 4.0b10 (Swofford 2002). All characters were equally weighted and gaps were treated as missing data. Trees were inferred using the heuristic search option with TBR branch swapping and 1000 random sequence additions. Max-trees were set to 5000, branches of zero length were collapsed and all parsimonious trees were saved. Clade robustness was assessed using a bootstrap (BT) analysis with 1,000 replicates (Felsenstein 1985). Descriptive tree statistics tree length (TL), consistency index (CI), retention index (RI), rescaled consistency index (RC), and homoplasy index (HI) were calculated for each Maximum Parsimonious Tree (MPT) generated. Sequences were also analyzed using Maximum Likelihood (ML) with RAxML-HPC2 through the Cipres Science Gateway (www.phylo.org; Miller et al. 2009). Branch support (BS) for ML analysis was determined by 1000 bootstrap replicates.

Table 1 Names, sample numbers and corresponding GenBank accession numbers of ITS sequences used in this study

Species name	Sample no.	GenBank accession no.	References
<i>Acanthofungus rimosus</i>	Wu 9601-1	MF043521	He et al. (2019)
<i>Acanthophysellum cerussatum</i>	He 2208	KX306874	He et al. (2019)
<i>Aleurobotrys botryosus</i>	He 2712	KX306877	He et al. (2019)
<i>Antrodia tanakae</i>	CLZhao 720	MG231457	This study
<i>Armillaria aotearoa</i>	NZFS 2425	NR151846	Hood & Ramsfield 2016
<i>Auricularia angiospermarum</i>	BJFC 017274	NR151847	Wu et al. 2015
<i>Auricularia villosula</i>	CLZhao 1296	MG231464	This study
<i>Bankera fuliginealba</i>	REB-285	JN135196	He et al. (2019)
<i>Boletinellus merulioides</i>	2630a	KM248952	He et al. (2019)
<i>Bondarzewia berkeleyi</i>	Dai 12759	KJ583202	He et al. (2019)
<i>Boreostereum radiatum</i>	RLG-9717-Sp	HM536085	Garcia-Sandoval et al. 2011
<i>Brunneoporus malicola</i>	CLZhao 1530	MG231451	This study
<i>Candelabrochaete langloisii</i>	FP-110343	KY948793	He et al. (2019)
<i>Ceraceomyces serpens</i>	HHB-15692-Sp	KP135031	He et al. (2019)
<i>Cinereomyces lindbladii</i>	CBS 290.71	MH860129	Vu et al. 2019
<i>Cinereomyces lindbladii</i>	CLZhao 1523	MG231489	This study
<i>Coniolepiota spongodes</i>	ECV-2010a	HM488756	He et al. (2019)
<i>Coprinellus curtus</i>	SZMC-NL-2339	FM878016	He et al. (2019)
<i>Coprinus comatus</i>	AFTOL-ID 626	AY854066	He et al. (2019)
<i>Corioloopsis polyzona</i>	Cui 11040	KR605824	He et al. (2019)
<i>Crepatura ellipsospora</i>	CLZhao 697	MK343695	Ma & Zhao 2019
<i>Cyclomyces lamellatus</i>	Cui 7629	JQ279603	He et al. (2019)
<i>Cylindrobasidium laeve</i>	CLZhao 767	MG231497	This study
<i>Daedalea quercina</i>	FFUI-4	MN596945	Direct Submission
<i>Daedaleopsis confragosa</i>	CLZhao 1481	MG231506	This study
<i>Efibula americana</i>	FP-102165	KP135016	He et al. (2019)
<i>Eichleriella alliciens</i>	HHB 7194	KX262120	He et al. (2019)
<i>Exidiopsis effusa</i>	OM 19136	KX262145	He et al. (2019)
<i>Flammulina velutipes</i>	AFTOL-ID 558	AY854073	He et al. (2019)
<i>Fomitiporia mediterranea</i>	AFTOL-ID 688	AY854080	He et al. (2019)
<i>Funalia gallica</i>	CLZhao 1306	MG231491	This study
<i>Funalia trogii</i>	CLZhao 1557	MG231874	This study
<i>Gloeophyllum sepium</i>	CLZhao 732	MG231532	This study
<i>Gloeophyllum trabeum</i>	1320	HM536094	He et al. (2019)
<i>Gloeoporus taxicola</i>	CLZhao 1441	MG231549	This study
<i>Grifola frondosa</i>	AFTOL-ID 701	AY854084	He et al. (2019)
<i>Gymnopus confluens</i>	ZRL 20151148	LT716054	He et al. (2019)
<i>Heliocybe sulcata</i>	IBUG 9930	HM536095	He et al. (2019)
<i>Heterobasidium annosum</i>	06129/6	KJ583211	He et al. (2019)
<i>Heterobasidium insulare</i>	CLZhao 2899	MK268944	This study
<i>Heterobasidium orientale</i>	CLZhao 696	MG231561	This study
<i>Hymenochaetopsis yasudae</i>	CLZhao 1422	MG231607	This study
<i>Hymenopellis radicata</i>	AFTOL-ID 561	DQ241780	He et al. (2019)
<i>Hyphoderma macaronesticum</i>	TFC Mic 15939	NR119817	Schoch et al. 2014
<i>Hyphoderma setigerum</i>	CBS 421.72	MH860512	Vu et al. 2019
<i>Hyphoderma transiens</i>	CLZhao 1365	MK404378	This study
<i>Hyphoderma variolosum</i>	CBS 735.91	MH862321	Vu et al. 2019
<i>Hyphodermella rosae</i>	FP-150552	KP134978	He et al. (2019)
<i>Inocutis dryophilus</i>	DLL 2012-001	KU139186	He et al. (2019)
<i>Irpex lacteus</i>	CLZhao 1258	MG231709	This study
<i>Junghuhnia crustacea</i>	X 262	JN710553	Miettinen et al. 2012
<i>Lacrymaria lacrymabunda</i>	CBS 211.31	MH855192	He et al. (2019)
<i>Laurilia sulcata</i>	CBS 365.49	MH856552	Vu et al. 2019
<i>Lenzitopsis daii</i>	Yuan 2959	JN169799	He et al. (2019)

Table 1 Continued.

Species name	Sample no.	GenBank accession no.	References
<i>Lycoperdon ericaeum</i>	ZRL 20151498	LT716030	He et al. (2019)
<i>Macrolepiota dolichaula</i>	xml 2013058	LT716021	He et al. (2019)
<i>Megalocystidium luridum</i>	CBS 106.71	MH860024	Vu et al. 2019
<i>Megasporoporiella subcavernulosa</i>	CLZhao 1438	MG231737	This study
<i>Melanogaster rivularis</i>	S 190	HQ714731	He et al. (2019)
<i>Meripilus giganteus</i>	FP-135344	KP135307	He et al. (2019)
<i>Meruliopsis albostramineus</i>	HHB-10729	KP135051	He et al. (2019)
<i>Microporus xanthopus</i>	CLZhao 1285	MG231749	This study
<i>Nigroporus vinosus</i>	KA17-0261	MN294801	Direct Submission
<i>Oligoporus farinosus</i>	CIEFAP 91	JX090117	Pildain & Rajchenberg 2013
<i>Omphalotus olearius</i>	AFTOL-ID 1718	DQ494681	He et al. (2019)
<i>Perenniporia hainaniana</i>	Cui 6364	JQ861743	He et al. (2019)
<i>Phaeophlebiopsis caribbeana</i>	HHB-6990	KP135415	He et al. (2019)
<i>Phanerochaete concrescens</i>	CLZhao 1430	MG231768	This study
<i>Phanerochaete sordida</i>	CLZhao 1459	MG231774	This study
<i>Phellinus ellipsoideus</i>	Cui 4270	JQ837948	He et al. (2019)
<i>Phlebiopsis crassa</i>	CLZhao 786	MG231791	This study
<i>Phylloporus pelletieri</i>	Pp 1	DQ534566	He et al. (2019)
<i>Pisolithus tinctorius</i>	AWW 219	EU718114	He et al. (2019)
<i>Polyozellus multiplex</i>	AFTOL-ID 677	DQ411528	He et al. (2019)
<i>Polyporus arcularius</i>	CLZhao 1338	MG231798	This study
<i>Porphyrellus porphyrosporus</i>	MB 97-023	DQ534563	He et al. (2019)
<i>Postia lactea</i>	Cui 9319	KX900894	Direct Submission
<i>Psathyrella candolleana</i>	ZRL 20151400	LT716063	He et al. (2019)
<i>Pseudochaete subrigidula</i>	He 1157	JQ716403	He et al. (2019)
<i>Pycnoporus sanguineus</i>	ZRL 2015009	LT716078	He et al. (2019)
<i>Rhizochaete americana</i>	FP-102188	KP135409	He et al. (2019)
<i>Rhizomarasmius oreinus</i>	AQUI 6763	NR132910	Moreau et al. 2015
<i>Rhodocollybia maculata</i>	AFTOL-ID 540	DQ404383	He et al. (2019)
<i>Rhodotus asperior</i>	HKAS 56754	KC179737	He et al. (2019)
<i>Rigidoporus undatus</i>	Miettinen 13591	KY948731	He et al. (2019)
<i>Sarcodon joeides</i>	REB-270	KC571772	He et al. (2019)
<i>Schizophyllum commune</i>	CLZhao 1528	MG231811	This study
<i>Schizophyllum leprieurii</i>	ROBLEDO 1313	KM098065	Direct Submission
<i>Scleroderma areolatum</i>	AWW 211	EU718115	He et al. (2019)
<i>Sparsitubus nelumbiformis</i>	Cui 8497	KX880631	He et al. (2019)
<i>Steccherinum bourdotii</i>	CLZhao 1347	MG231820	This study
<i>Steccherinum ochraceum</i>	CLZhao 2897	MK269280	This study
<i>Stereum hirsutum</i>	CLZhao 1411	MG231830	This study
<i>Stereum rugosum</i>	CLZhao 1310	MG231836	This study
<i>Stereum sanguinolentum</i>	CLZhao 668	MG231838	This study
<i>Trametes hirsuta</i>	CLZhao 1544	MG231868	This study
<i>Trametopsis cervina</i>	TJV 93 216T	JN165020	He et al. (2019)
<i>Tremella flava</i>	CBS 8471	KY105681	He et al. (2019)
<i>Tremella mesenterica</i>	CBS 6973	NR155937	He et al. (2019)
<i>Trulla dentipora</i>	AS 2288	KY970064	Direct Submission
<i>Veluticeps fimbriata</i>	L-10628-Sp	HM536100	He et al. (2019)
<i>Xylobolus frustulatus</i>	He 2231	KU881905	He et al. (2019)

MrModeltest 2.3 (Nylander 2004) was used to determine the best-fit evolution model for each data set for Bayesian inference (BI). Bayesian inference was calculated with MrBayes3.1.2 with a general time reversible (GTR+I+G) model of DNA substitution and a gamma distribution rate variation across sites (Ronquist & Huelsenbeck 2003). Four Markov chains were run for 2 runs

from random starting trees for 1500 thousand generations (Fig. 1), and trees were sampled every 100 generations. The first one-fourth generations were discarded as burn-in. A majority rule consensus tree of all remaining trees was calculated. A majority rule consensus tree of all remaining trees was calculated. Branches were considered as significantly supported if they received maximum likelihood bootstrap (BS) >70%, maximum parsimony bootstrap (BT) >50%, or Bayesian posterior probabilities (BPP) >0.95.

Results

Molecular phylogeny

The ITS dataset (Fig. 1) included sequences from 102 fungal specimens representing 101 species. The dataset had an aligned length of 1389 characters, of which 308 characters were constant, 283 parsimony-uninformative, and 798 parsimony-informative. Maximum parsimony analysis yielded 1 equally parsimonious tree (TL = 8644, CI = 0.2678, HI = 0.7322, RI = 0.4535, RC = 0.1215). The best-fit model for ITS alignment estimated and applied in the Bayesian was GTR+I+G, lset nst = 6, rates = invgamma; prset statefreqpr = dirichlet (1,1,1,1). Bayesian resulted in a similar topology with an average standard deviation of split frequencies = 0.029534 (BI), and the effective sample size (ESS) across the two runs is the double of the average ESS (avg ESS) = 354.

The phylogeny (Fig. 1) inferred from ITS sequences demonstrated that fifty-two species nested in sixteen families, Auriculariaceae, Bondarzewiaceae, Dacrybolaceae, Fomitopsidaceae, Gelatoporiaceae, Gloeophyllaceae, Hymenochaetaceae, Hyphodermataceae, Irpicaceae, Phanerochaetaceae, Physalacriaceae, Polyporaceae, Schizophyllaceae, Schizoporaceae, Steccherinaceae and Stereaceae, belonging to six orders Agaricales, Auriculariales, Gloeophyllales, Hymenochaetales, Polyporales, Russulales in Agaricomycetes.

Checklist

An alphabetical list (according to genus name) of wood-decaying fungi identified in these investigations is given below. The authors of scientific names are according to the second edition of Authors of Fungal Names (<http://www.indexfungorum.org/AuthorsOfFungalNames.html>). Substrate and collecting data are provided after the name of each species. The hosts are listed alphabetically, and within the same host tree, they are arranged by the order: living tree, dead standing tree, trunk, fallen branch and stump. The collectors and collection numbers are listed alphabetically, too (Dai 2011, 2012a).

1. *Antrodia tanakae* (Murrill) Spirin & Miettinen, on the fallen branch of *Acacia dealbata* Link, CLZhao 720; on the stump of *Acacia dealbata*, CLZhao 1536
2. *Auricularia villosula* Malysheva, on the fallen branch of *Alnus nepalensis*, CLZhao 1428; on the trunk of *Juglans regia* L., CLZhao 743; on the trunk of *Quercus*, CLZhao 1296; on the fallen branch of *Quercus*, CLZhao 1340
3. *Basidioradulum crustosum* (Pers.) Zmitr., Malysheva & Spirin, on the fallen angiosperm branch, CLZhao 3028
4. *Bjerkandera adusta* (Willd.) P. Karst, on the stump of *Pinus yunnanensis*, CLZhao 1555; on the dead tree of *Quercus*, CLZhao 1275
5. *Brunneoporus malicola* (Berk. & M.A. Curtis) Audet, on the stump of *Acacia dealbata*, CLZhao 1524; on the trunk of *Quercus acutissima* Carr., CLZhao 1530
6. *Byssomerulius corium* (Pers.) Parmasto, on the fallen angiosperm branch, CLZhao 1560; on the fallen branch of *Pinus yunnanensis*, CLZhao 734, CLZhao 781; on the fallen branch of *Quercus*, CLZhao 693, CLZhao 1266, CLZhao 1274, CLZhao 1313; on the stump of *Quercus*, CLZhao 1341
7. *Cinereomyces lindbladii* (Berk.) Jülich, on the trunk of *Pinus yunnanensis*, CLZhao 1523
8. *Crepatura ellipsospora* C.L. Zhao, on the trunk of *Alnus*, CLZhao 697; on the fallen branch of *Quercus*, CLZhao 868, CLZhao 1260, CLZhao 1265

9. *Cylindrobasidium laeve* (Pers.) Chamuris, on the dead bamboo, CLZhao 756, CLZhao 767
10. *Daedaleopsis confragosa* (Bolton) J. Schröt, on the trunk of *Alnus nepalensis*, CLZhao 1481
11. *Funalia gallica* (Fr.) Bondartsev & Singer, on the trunk of *Alnus nepalensis*, CLZhao 1309; on the fallen branch of *Alnus nepalensis*, CLZhao 1306
12. *Funalia trogii* Berk., on the trunk of *Alnus nepalensis*, CLZhao 1540; on the angiosperm trunk, CLZhao 741, CLZhao 3009, CLZhao 1557; on the fallen angiosperm branch CLZhao 1552
13. *Fuscoporia torulosa* (Pers.) T. Wagner & M. Fisch, on the fallen branch of *Quercus*, CLZhao 1305
14. *Gloeophyllum sepiarium* (Wulfen) P. Karst, on the trunk of *Pinus yunnanensis*, CLZhao 732, CLZhao 764; on the fallen branch of *Pinus yunnanensis*, CLZhao 774, CLZhao 904; on the stump of *Pinus yunnanensis*, CLZhao 731, CLZhao 784
15. *Gloeoporus dichrous* (Fr.) Bres., on the fallen branch of *Alnus nepalensis*, CLZhao 1471
16. *Gloeoporus taxicola* (Pers.) Gilb. & Ryvardeen, on the trunk of *Pinus armandii*, CLZhao 1441
17. *Heterobasidium insulare* (Murrill) Ryvardeen, on the stump of *Pinus yunnanensis*, CLZhao 2899
18. *Heterobasidium orientale* Tokuda, T. Hatt. & Y.C. Dai, trunk of *Pinus yunnanensis*, CLZhao 696
19. *Hymenochaete adusta* (Lév.) Har. & Pat., J. Bot, on the trunk of *Alnus*, CLZhao 700
20. *Hymenochaete rheicolor* (Mont.) Lév., on the trunk of *Alnus*, CLZhao 672, CLZhao 679; on the fallen branch of *Quercus*, CLZhao 666, CLZhao 671, CLZhao 682
21. *Hymenochaete villosa* (Lév.) Bres., on the trunk of *Quercus acutissima*, CLZhao 1533
22. *Hymenochaetopsis corrugata* (Fr.) S.H. He & Jiao Yang, on the fallen angiosperm branch, CLZhao 2893
23. *Hymenochaetopsis yasudae* (Imazeki) S.H. He & Jiao Yang, on the fallen branch of *Alnus nepalensis*, CLZhao 1422, CLZhao 1445; on the living tree of *Pinus armandii*, CLZhao 1475; on the fallen branch of *Pinus armandii*, CLZhao 1486, CLZhao 1495, CLZhao 1549
24. *Hyphoderma transiens* (Bres.) Parmasto, on the fallen angiosperm branch, CLZhao 1493; on the trunk of *Populus yunnanensis* Dode, CLZhao 1365
25. *Hyphodontia tropica* Sheng H. Wu, on the fallen angiosperm branch, CLZhao 2898, CLZhao 2901
26. *Irpex lacteus* (Fr.) Fr., on the trunk of *Acacia dealbata*, CLZhao 1258
27. *Lopharia cinerascens* (Schwein.) G. Cunn., on the fallen branch of *Alnus nepalensis*, CLZhao 1499
28. *Megasporoporiella subcavernulosa* (Y.C. Dai & Sheng H. Wu) B.K. Cui, on the fallen branch of *Alnus nepalensis*, CLZhao 1438, CLZhao 1466; on the stump of *Alnus nepalensis*, CLZhao 1412; on the fallen angiosperm branch, CLZhao 2966, CLZhao 2984, CLZhao 3016; on the stump of *Cupressus funebris* Endl., CLZhao 1444, CLZhao 1491
29. *Microporus xanthopus* (Fr.) Kuntze, on the fallen branch of *Alnus nepalensis*, CLZhao 1503; on the fallen angiosperm branch, CLZhao 3012; on the trunk of *Quercus*, CLZhao 1253, CLZhao 1285; on the fallen branch of *Quercus*, CLZhao 1268, CLZhao 1304; on the stump of *Quercus*, CLZhao 1343
30. *Phanerochaete concrescens* Spirin & Volobuev, on the fallen branch of *Alnus nepalensis*, CLZhao 1541, CLZhao 1545; on the fallen angiosperm branch, CLZhao 2929, CLZhao 2931, CLZhao 2939, CLZhao 2940, CLZhao 2945, CLZhao 2946, CLZhao 2949; on the trunk of *Pinus yunnanensis*, CLZhao 2916
31. *Phanerochaete sordida* (P. Karst.) J. Erikss. & Ryvardeen, on the trunk of *Alnus*, CLZhao 698; on the fallen branch of *Alnus nepalensis*, CLZhao 1459, CLZhao 1541, CLZhao 1545; on the angiosperm trunk, CLZhao 1461; on the fallen angiosperm branch, CLZhao 2929, CLZhao 2931, CLZhao 2939, CLZhao 2940, CLZhao 2945, CLZhao 2946, CLZhao 2949, CLZhao 4738, CLZhao 4746, CLZhao 4754; on the fallen branch of *Pinus armandii*, CLZhao 1515; on the trunk of *Pinus yunnanensis*, CLZhao 2916
32. *Phellinus gilvus* (Schwein.) Pat., on the fallen branch of *Alnus nepalensis*, CLZhao 1334

33. *Phlebiopsis crassa* (Lév.) Floudas & Hibbett, on the angiosperm trunk, CLZhao 786; on the fallen branch of *Coriaria nepalensis* Wall., CLZhao 1295, CLZhao 1308; on the trunk of *Pinus yunnanensis*, CLZhao 724; on the trunk of *Quercus*, CLZhao 1269, CLZhao 1314; on the fallen branch of *Quercus acutissima*, CLZhao 1532
34. *Polyporus arcularius* (Batsch) Fr. on the stump of *Coriaria nepalensis*, CLZhao 1338; on the fallen branch of *Quercus*, CLZhao 1316
35. *Postia hibernica* (Berk. Broome) Jülich, on the fallen angiosperm branch, CLZhao 2903; on the fallen branch of *Pinus yunnanensis*, CLZhao 2909
36. *Pulcherricium coeruleum* (Lam.) Parmasto, on the fallen branch of *Alnus nepalensis*, CLZhao 1434; on the fallen angiosperm branch, CLZhao 3003, CLZhao 3020
37. *Schizophyllum commune* Fr., on the trunk of *Acacia dealbata*, CLZhao 1527; on the fallen branch of *Acacia dealbata*, CLZhao 1528, CLZhao 1529; on the trunk of *Alnus nepalensis*, CLZhao 1537; on the stump of *Alnus nepalensis*, CLZhao 1562; on the stump of *Eucalyptus robusta*, CLZhao 1561
38. *Steccherinum bourdotii* Saliba & A. David, on the fallen branch of *Alnus nepalensis*, CLZhao 1472; on the fallen branch of *Pinus armandii*, CLZhao 1347
39. *Steccherinum ochraceum* (Pers. ex J.F. Gmel.) Gray, on the fallen angiosperm branch, CLZhao 2897, CLZhao 2968
40. *Stereum gausapatum* (Fr.) Fr., on the trunk of *Alder*, 11 January 2017, CLZhao 668, CLZhao 669, CLZhao 673, CLZhao 677, CLZhao 683, CLZhao 694, CLZhao 707; on the dead tree of *Quercus*, 22 April 2017, CLZhao 1259, CLZhao 1270, CLZhao 1290, CLZhao 1300, CLZhao 1318, CLZhao 1320
41. *Stereum hirsutum* (Willd.) Pers., on the fallen branch of *Alnus nepalensis*, CLZhao 1404, CLZhao 1405, CLZhao 1427, CLZhao 1455, CLZhao 1457, CLZhao 1465, CLZhao 1469, CLZhao 1470, CLZhao 1479, CLZhao 1482, CLZhao 1489, CLZhao 1498; on the fallen angiosperm branch, CLZhao 740; on fallen branch of *Coriaria nepalensis*, CLZhao 1559; on fallen branch of *Pinus yunnanensis*, CLZhao 2906; on the fallen branch of *Quercus*, CLZhao 1291
42. *Stereum rugosum* Pers., on the trunk of *Quercus*, CLZhao 1310
43. *Stereum sanguinolentum* (Alb. & Schwein.) Fr., on the trunk of *Alnus*, CLZhao 673; on the stump of *Pinus yunnanensis*, CLZhao 669
44. *Trametes hirsuta* (Wulfen) Lloyd, on the fallen branch of *Alnus nepalensis*, CLZhao 1544; on the angiosperm trunk, CLZhao 1344; on the trunk of *Pinus yunnanensis*, CLZhao 739
45. *Trametes strumosa* (Fr.) Zmitr., on the angiosperm trunk, CLZhao 718
46. *Trametes versicolor* (L.) Lloyd, on the trunk of *Alnus nepalensis*, CLZhao 1539, CLZhao 1546; on the fallen branch of *Alnus nepalensis*, CLZhao 1431; on the stump of *Alnus nepalensis*, CLZhao 1510; on the angiosperm trunk, CLZhao 748, CLZhao 1293, CLZhao 1477; on the trunk of *Cupressus funebris*, CLZhao 1509; on the trunk of *Quercus*, CLZhao 1330; on the fallen branch of *Quercus*, CLZhao 686, CLZhao 714, CLZhao 1302, CLZhao 1307
47. *Trametopsis cervina* (Schwein.) Tomšovský, on the trunk of *Alnus nepalensis*, CLZhao 1246; on the fallen branch of *Quercus*, CLZhao 1315
48. *Trichaptum abietinum* (Dicks.) Ryvar den, on the angiosperm stump, CLZhao 719, CLZhao 723, CLZhao 736, CLZhao 777, CLZhao 3004; on the trunk of *Pinus yunnanensis*, CLZhao 730
49. *Tubulicrinis xantha* C.L. Zhao, on the fallen branch of *Pinus yunnanensis*, CLZhao 2868, CLZhao 2869, CLZhao 2883
50. *Xylodon kunmingensis* L. W. Zhou & C.L. Zhao, on the fallen angiosperm branch, CLZhao 3010, CLZhao 3019; on the stump of angiosperm, CLZhao 755; on the fallen branch of *Pinus yunnanensis*, CLZhao 752
51. *Xylodon nespori* (Bres.) Hjortstam & Ryvar den, on the trunk of *Cupressus funebris*, CLZhao 1492
52. *Xylodon rimosissimus* (Peck) Hjortstam & Ryvar den, on the fallen branch of *Pinus armandii*, CLZhao 1487




Fig. 1 – Maximum parsimony strict consensus tree illustrating the phylogeny of 52 species with related taxa in Agaricomycetes based on ITS sequences. Branches are labelled with maximum likelihood bootstrap equal to or greater than 70%, parsimony bootstrap equal to or greater than 50% and Bayesian posterior probabilities equal to or greater than 0.97, respectively. The taxa from the present study are indicated in black bold.




Fig. 1 – Continued.

Discussion

In the kingdom Fungi, the phyla Ascomycota and Basidiomycota cover around 97% of all fungal species (Willis 2018). According to the latest version of Ainsworth & Bisby's Dictionary of the Fungi (Kirk et al. 2008), there are 1589 genera and more than 30 thousand species of Basidiomycota, which comprise nearly 32% of all described fungal taxa (Dai et al. 2015), and more and more taxa were recorded worldwide every year, which observably increase the fungal diversity (Sarma & Hyde 2018, Yafetto 2018, Freitas-Neto et al. 2019, Hipol et al. 2019, Ayesha et al. 2020, De Leon AM et al. 2020, Huang et al. 2020, Himani & Krishnappa 2020, Blanco-Dios 2021, Kumar et al. 2021, Wang & Zhao 2021, Zong et al. 2021).

Although some notable explorations of wood-decaying fungi have been made from Yunnan Province (Cui & Dai 2012, Wu et al. 2017, Zhao & Wu 2017, Shen et al. 2018, Wu et al. 2018, Liu et al. 2019, Luo et al. 2019, Xu et al. 2019, Chen & Zhao 2020, Huang et al. 2020, Wang et al. 2020, Wu et al. 2021, Zong et al. 2021), only three new species were found from Haikou Forestry Farm (Ma & Zhao 2019, Shi et al. 2019, He et al. 2020). Species in the present list are mostly new to the studied area. The 681 specimens belong to 52 species, which are distributed in 37 genera, 16 families, 6 orders, in which 11 are pathogenic (Dai 2012b), and 28 are medicinal (Dai & Yang 2008), and a few are edible (Dai et al. 2010). Among them, 28 species belong to order Polyporales and 14 species belong to order Hymenochaetales and 6 species belong to order Russulales, and 2 species belong to order Agaricales and 1 species belong to order Auriculariales and 1 species belong to order Gloeophyllales.

The molecular phylogenetic analyses with combined nLSU, SSU, 5.8s, rpb1, rpb2, and tef1 datasets for the subphyla Agaricomycotina, Pucciniomycotina and Ustilaginomycotina revealed that 1928 currently used genera names are distributed in 241 families, 68 orders, 18 classes and four subphyla (He et al. 2019). In the present study, 52 species nested in 37 genera, 16 families, 6 orders based on ITS dataset, which is similar to the topology of the phylogenetic analyses in He et al. (2019).

This work will comprehensively improve the understanding of the diversity of wood-decaying fungi in this area, which is conducive to the rational utilization and effective protection of fungal resources, and provides scientific basis for the prevention and control of forest diseases in this farm.

Acknowledgements

The researches were supported by the Yunnan Fundamental Research Project (Grant No. 202001AS070043) and the Science Foundation of Southwest Forestry University (Project No. 111715).

References

- Ayesha P, Maham A, Ariza A, Husnain H. 2020 – “*Ganoderma lucidum sensu lato*” – a sacred mushroom for immortality. *Studies in Fungi* 5(1), 508–516. Doi 10.5943/sif/5/1/31
- Blanco-Dios JB. 2021 – Notes on the genus *Entoloma* s.l. in the Northwest of the Iberian Peninsula (XI): a new species in subgenus *Entoloma*. *Studies in Fungi* 6(1), 64–70. Doi 10.5943/sif/6/1/2
- Chen JZ, Zhao CL. 2020 – Morphological and molecular identification of four new resupinate species of *Lyomyces* (Hymenochaetales) from southern China. *Mycosystema* 65, 101–118. Doi 10.3897/mycokeys.65.48660
- Cui BK, Dai YC. 2012 – Wood-decaying fungi in eastern Himalayas 3. Polypores from Laojunshan Mountains, Yunnan Province. *Mycosystema* 31(4), 486–492.
- Dai YC. 2011 – A revised checklist of corticioid and hydroid fungi in China for 2010. *Mycoscience* 52, 69–79. Doi: 10.1007/S10267-010-0068-1
- Dai YC. 2012a – Polypore diversity in China with an annotated checklist of Chinese polypores. *Mycoscience* 53, 49–80. Doi 10.1007/s10267-011-0134-3
- Dai YC. 2012b – Pathogenic wood-decaying fungi on woody plants in China. *Mycosystema* 31(4), 493–509.
- Dai YC, Cui BK, Si J, He SH et al. 2015 – Dynamics of the worldwide number of fungi with emphasis on fungal diversity in China. *Mycological Progress* 14, 62. Doi 10.1007/s11557-015-1084-5
- Dai YC, Yang ZL. 2008 – A revised checklist of medicinal fungi in China. *Mycosystema* 27(6), 801–824. Doi 10.3969/j.issn.1672-6472.2008.06.001
- Dai YC, Zhou LW, Yang ZL, Wen HA et al. 2010 – A revised checklist of edible fungi in China. *Mycosystema* 29(1), 1–21.

- De Leon AM, Dulay AR, Villanueva AL, Kalaw SP. 2020 – Optimal culture conditions and toxicity assessment of *Fomitopsis feei* (Fr.): a newly documented macro fungus from Philippines. *Studies in Fungi* 5(1), 491–507. Doi 10.5943/sif/5/1/30
- Felsenstein J. 1985 – Confidence intervals on phylogenetics: an approach using bootstrap. *Evolution* 39, 783–791. Doi 10.1111/j.1558-5646.1985.tb00420.x
- Freitas-Neto JF, Sousa JO, Ovrebo CL, Baseia IG. 2019 – *Geastrum echinulatum* and *G. rusticum* (Geastraceae, Basidiomycota) – two new records for Central America. *Studies in Fungi* 4(1), 14–20. Doi 10.5943/sif/4/1/2
- Garcia-Sandoval R, Wang Z, Binder M, Hibbett DS. 2011 – Molecular phylogenetics of the Gloeophyllales and relative ages of clades of Agaricomycotina producing a brown rot. *Mycologia* 103(3), 510–524. Doi 10.3852/10-209
- Hall TA. 1999 – Bioedit: a user-friendly biological sequence alignment editor and analysis program for Windows 95/98/NT. *Nucleic Acids Symposium Series* 41, 95–98.
- He MQ, Zhao RL, Hyde KD, Begerow D et al. 2019 – Notes, outline and divergence times of Basidiomycota. *Fungal Diversity* 99(1), 105–367. Doi 10.1007/s13225-019-00435-4
- He X, Shi ZJ, Zhao CL. 2020 – Morphological and molecular identification of two new species of *Tubulicrinis* (Hymenochaetaceae, Hymenochaetales) from southern china. *Mycoscience* 61(4), 184–189. Doi 10.1016/j.myc.2020.03.008
- Himani S, Krishnappa M. 2020 – *Xylaria oxyacanthae* (Xylariaceae), a new record on *Diospyros melanoxylon* from India. *Studies in Fungi* 5(1), 485–490. Doi 10.5943/sif/5/1/29
- Hipol RM, Baldelomar JA, Bolinget KC, Solis AFF. 2019 – The soil fungi producing siderophores of Mt. Yangbew, Tawang, La Trinidad, Benguet. *Studies in Fungi* 4(1), 1–13. Doi 10.5943/sif/4/1/1
- Hood IA, Ramsfield TD. 2016 – *Armillaria aotearoa* species nova. *New Zealand Journal of Forestry Science* 46(1), 2. Doi 10.1186/s40490-016-0058-y
- Huang RX, Luo KY, Zhao CL. 2020 – *Phlebia nigrodontea* sp. nov. in Meruliaceae (Polyporales) with a black hymenial surface. *Phytotaxa* 458(3), 195–206. Doi 10.11646/phytotaxa.458.3.2.
- Kirk PM, Cannon PF, Minter DW, Stalpers JA. 2008 – *Ainsworth & Bisby's Dictionary of the Fungi*, 10th edn. CAB International, Wallingford.
- Kumar S, Singh R, Kamal. 2021 – Global diversity and distribution of distoseptosporic micromycete *Corynespora* Güssow (Corynesporascaceae): An updated checklist with current status. *Studies in Fungi* 6(1), 1–63. Doi: 10.5943/sif/6/1/1
- Liu XF, Shen S, Zhao CL. 2019 – Morphological and molecular identification of a new species of *Eichleriella* (Auriculariales, Basidiomycota) in China. *Phytotaxa* 404(6), 245–254. Doi 10.11646/phytotaxa.404.6.3
- Luo KY, Ma X, Zhao CL. 2019 – *Neofavolus yunnanensis* sp. nov. (Polyporales, Basidiomycota) from China: evidence from morphology and DNA sequence data. *Phytotaxa* 408, 109–116. Doi 10.11646/phytotaxa.408.2.3
- Ma X, Zhao CL. 2019 – *Crepatura ellipsospora* gen. et sp. nov. in Phanerochaetaceae (Polyporales, Basidiomycota) bearing a tuberculate hymenial surface. *Mycological Progress* 18(6), 785–793. Doi 10.1007/s11557-019-01488-0
- M'Barek HN, Arif S, Taidi B, Hajjaj H. 2020 – Consolidated bioethanol production from olive mill waste: Wood-decay fungi from central Morocco as promising decomposition and fermentation biocatalysts. *Biotechnology Reports*, 28. Doi 10.1016/j.btre.2020.e00541
- Miettinen O, Larsson E, Sjökvist E, Larsson KH. 2012 – Comprehensive taxon sampling reveals unaccounted diversity and morphological plasticity in a group of dimitic polypores (Polyporales, Basidiomycota). *Cladistics-the International Journal of the Willi Hennig Society* 28(3), 251–270. Doi 10.1111/j.1096-0031.2011.00380.x
- Miller MA, Holder MT, Vos R, Midford PE et al. 2009 – The CIPRES Portals. CIPRES. http://www.phylo.org/sub_sections/portal. 2009-08-04. (Archived by WebCite at <http://www.webcitation.org/5imQlJeQa>).

- Moreau PA, Vila J, Aime MC, Antonin V et al. 2015 – *Cibaomyces* and *Cyptotrampa*, two new genera for Europe, and an emendation of *Rhizomarasmius* (Basidiomycota, Physalacriaceae). *Mycological Progress* 14, 4. Doi 10.1007/s11557-015-1024-4
- Nylander JAA. 2004 – MrModeltest v2. Program distributed by the author. Evolutionary Biology Centre, Uppsala University.
- Petersen JH. 1996 – Farvekort. The Danish Mycological Society's colour-chart. Foreningen til Svampekundskabens Fremme, Greve.
- Pildain MB, Rajchenberg M. 2013 – The phylogenetic position of *Postia* s.l. (Polyporales, Basidiomycota) from Patagonia, Argentina. *Mycologia* 105(2), 357–367. Doi 10.3852/12-088
- Ronquist F, Huelsenbeck JP. 2003 – MrBayes 3: Bayesian phylogenetic inference under mixed models. *Bioinformatics* 19, 1572–1574. Doi 10.1093/bioinformatics/btg180
- Runnel K, Miettinen O, Lohmus A. 2021 – Polypore fungi as a flagship group to indicate changes in biodiversity – a test case from Estonia. *IMA Fungus* 12(1). Doi 10.1186/s43008-020-00050-y
- Russell R, Paterson M. 2006 – *Ganoderma* - A therapeutic fungal biofactory. *Phytochemistry* 67(18), 1985–2001. Doi 10.1016/j.phytochem.2006.07.004
- Sarma VV, Hyde KD. 2018 – Fungal species consortia on *Nypa fruticans* at Brunei. *Studies in Fungi* 3(1), 19–26. Doi 10.5943/sif/3/1/3
- Shen S, Ma X, Xu TM, Zhao CL. 2018 – *Phlebia ailaoshanensis* sp. nov. (Polyporales, Basidiomycota) evidenced by morphological characters and phylogenetic analyses. *Phytotaxa* 373, 184–196. Doi 10.11646/phytotaxa.373.3.2
- Shi ZW, Wang XW, Zhou LW, Zhao CL. 2019 – *Xylodon kunmingensis* sp. nov. (Hymenochaetales, Basidiomycota) from southern China. *Mycoscience* 60, 184–188. Doi 10.1016/j.myc.2019.02.002
- Schoch CL, Robbertse B, Robert V, Vu D et al. 2014 – Finding needles in haystacks: linking scientific names, reference specimens and molecular data for Fungi. *Database*, 1–21. Doi 10.1093/database/bau061
- Swofford DL. 2002 – PAUP*: phylogenetic analysis using parsimony (*and other methods). Version 4.0b10. Sinauer Associates, Massachusetts.
- Vinay DS, Ryan EP, Pawelec G, Talib WH et al. 2015 – Immune evasion in cancer: Mechanistic basis and therapeutic strategies. *Seminars in Cancer Biology* 35, S185–S198. Doi 10.1016/j.semcancer.2015.03.004
- Vu D, Groenewald M, De VM, Gehrman T et al. 2019 – Large-scale generation and analysis of filamentous fungal DNA barcodes boosts coverage for kingdom fungi and reveals thresholds for fungal species and higher taxon delimitation. *Studies in Mycology* 92, 135–154. Doi 10.1016/j.simyco.2018.05.001
- Wang RZ, Yuan ZY, Han CZ, Zhao CL. 2020 – Morphological and molecular identification of a new species of *Cinereomyces* (Polyporales, Basidiomycota) in southern China. *Phytotaxa* 459(1), 051–060. Doi 10.11646/phytotaxa.459.1.5
- Wang H, Zhao CL. 2021 – *Hyphodermella aurantiaca* sp. nova. (Polyporales, Basidiomycota) as evidenced by morphological characters and phylogenetic analyses. *Annales Botanici Fennici* 58, 61–68.
- White TJ, Bruns T, Lee S, Taylor J. 1990 – Amplification and direct sequencing of fungal ribosomal RNA genes for phylogenetics. In: Innis MA, Gelfand DH, Sninsky JJ, White TJ (eds), *PCR Protocols: a guide to methods and applications*. Academic Press, New York, pp. 315–322. Doi 10.1016/B978-0-12-372180-8.50042-1
- Willis KJ. (ed.) 2018 – State of the World's Fungi 2018. Report. Royal Botanic Gardens, Kew.
- Wu F, Yuan Y, He SH, Bandaraet AR et al. 2015 – Global diversity and taxonomy of the *Auricularia auricula-judae* complex (Auriculariales, Basidiomycota). *Mycological Progress* 14, 95. Doi 10.1007/s11557-015-1113-4
- Wu SH, Chen CL, Wei CL, Chen YP et al. 2020 – *Ganoderma bambusicola* sp. nov. (Polyporales, Basidiomycota) from southern Asia. *Phytotaxa* 456(1), 75–85.

Doi 10.11646/phytotaxa.456.1.5

- Wu YX, Wu JR, Zhao CL. 2021 – *Steccherinum tenuissimum* and *S. xanthum* spp. nov. (Polyporales, Basidiomycota): New species from China. PLoS One 16(1), e0244520. Doi 10.1371/journal.pone.0244520
- Wu ZQ, Shen S, Luo KY, Wang ZH et al. 2017 – Morphological and molecular identification of a new species of *Atraporrella* (Polyporales, Basidiomycota) in China. Phytotaxa 332, 31–40. Doi 10.11646/phytotaxa.332.1.3
- Wu ZQ, Xu TM, Shen S, Liu XF et al. 2018 – *Elaphroporia ailaoshanensis* gen. et sp. nov. in Polyporales (Basidiomycota). MycoKeys 29, 81–95. Doi 10.3897/mycokeys.29.22086
- Xu SF, Li XY, Xia J, Liao XK. 2015 – Evaluation on forest health condition in Haikou Forest Farm of Kunming. Journal of Southwest Forestry University 35(2), 68–72.
- Xu TM, Liu XF, Chen YH, Zhao CL. 2019 – *Rhomboidia wuliangshanensis* gen. et sp. nov. from southwestern China. Mycotaxon 134, 649–662. Doi 10.5248/134.649
- Xiong XW, Zhou HQ. 2019 – Evaluation of health status of *Pinus armandii* plantation in central Yunnan plateau: a case study of state-owned Haikou Forest Farm in Kunming. Journal of Jiangsu Forestry Science & Technology 46(3), 1–5.
- Yafetto L. 2018 – Protein enrichment of cassava pulp by solid-state fermentation using *Aspergillus niger*. Studies in Fungi 3(1), 7–18. Doi 10.5943/sif/3/1/2
- Zhang ZH, Wang YT, Tie XR, Chen JL. 2017 – The effectiveness evaluation about natural forest protecti on project construction in Kunming Haikou Forest Farm. Journal of Green Science and Technology 9, 142–146.
- Zhao CL, Wu ZQ. 2017 – *Ceriporiopsis kunmingensis* sp. nov. (Polyporales, Basidiomycota) evidenced by morphological characters and phylogenetic analysis. Mycological Progress 16, 93–100. Doi 10.1007/s11557-016-1259-8
- Zong TK, Liu CM, Wu JR, Zhao CL. 2021 – *Trechispora daweishanensis* and *T. xantha* spp. nov. (Hydnodontaceae, Trechisporales) found in Yunnan Province of China. Phytotaxa 479(2), 147–159. Doi 10.11646/phytotaxa.479.2.1